

## Walter Keller

### List of publications

#### Original research reports:

Stüttgen, G., W. Keller, and K. Knobel: Zum Einfluss von Heparinen und Adenosinphosphorsäuren auf Substrate und Aktivität der Mono- und Diaminoxidasen. *Z. ges. exp. Med.* 134, 446-543 (1961).

Keller, W., O. von Deimling, R. Rossner, and W. Oehlert: Hormonabhängige Enzymverteilung in Geweben: XI. Geschlechts- und zyklusbedingte Unterschiede des Einbaus von <sup>3</sup>H-Cytidin in die Hauptstückepithelien der Rattenniere. *Histochemie* 12, 159-164 (1968).

von Deimling, O. and W. Keller: Hormonabhängige Enzymverteilung in Geweben: XII. Der spezifische Einbau von <sup>3</sup>H-Cytidin in die Kerne der Hauptstückepithelien der Rattenniere unter Oestradiol- und Actinomycineinwirkung. *Histochemie* 15, 333-347 (1968).

Goedde, H.W., U. Langenbeck, D. Brackertz, W. Keller, T. Rokkones, S. Halvorsen, R. Kiil, and B. Merton: Clinical and biochemical-genetic aspects of intermittent branched-chain ketoaciduria. *Acta Paediat. Scand.* 59, 83-87 (1970).

Keller, W. and R. Goor: Mammalian RNA polymerase. Structural and functional properties. *Cold Spring Harbor Symp. Quant. Biol.* 35, 671-680 (1970).

Langenbeck, U., H.W. Rüdiger, M. Schulze-Schenking, W. Keller, D. Brackertz, and H.W. Goedde: Evaluation of a heterozygote test for maple syrup urine disease in leukocytes and cultured fibroblasts. *Humangenetik* 11, 304-315 (1971).

Keller, W.: RNA-primed DNA synthesis *in vitro*. *Proc. Nat. Acad. Sci. USA* 69, 1560-1564 (1972).

Sambrook, J., P.A. Sharp, and W. Keller: Transcription of Simian Virus 40. I. Separation of the strands of SV40 DNA and hybridization of the separated strands to RNA extracted from lytically infected and transformed cells. *J. Mol. Biol.* 70, 57-71 (1972).

Keller, W. and R. Crouch: Degradation of DNA-RNA hybrids by ribonuclease H and DNA polymerases of cellular and viral origin. *Proc. Nat. Acad. Sci. USA* 69, 3360-3364 (1972).

- Sugden, B. and W. Keller: Mammalian DNA-dependent ribonucleic acid polymerases. I. Purification and properties of an  $\alpha$ -amanitin-sensitive RNA polymerase and stimulatory factors from HeLa and KB cells. *J. Biol. Chem.* 248, 3777-3788 (1973).
- Sambrook, J., B. Sugden, W. Keller, and P.A. Sharp: Transcription of Simian Virus 40. III. Mapping of "early" and "late" species of RNA. *Proc. Nat. Acad. Sci. USA* 70, 3711-3715 (1973).
- Roberts, R.J., J.R. Arrand, and W. Keller: The length of the terminal repetition in Adenovirus-2 DNA. *Proc. Nat. Acad. Sci. USA* 71, 3829-3833 (1974).
- Arrand, J.R., W. Keller, and R. Roberts: The extent of the terminal repetition in Adenovirus-2 DNA. *Cold Spring Harbor Symp. Quant. Biol.* 39, 401-407 (1975).
- Keller, W. and I. Wendel: Stepwise relaxation of supercoiled SV40 DNA. *Cold Spring Harbor Symp. Quant. Biol.* 39, 199-208 (1975).
- Mulder, C., J.R. Arrand, H. Delius, W. Keller, U. Pettersson, R.J. Roberts, and P.A. Sharp: Cleavage maps of DNA from Adenovirus types 2 and 5 by restriction endonucleases *Eco* RI and *Hpa* I. *Cold Spring Harbor Symp. Quant. Biol.* 39, 397-400 (1975).
- Keller, W.: Characterization of purified DNA-relaxing enzyme from human tissue culture cells. *Proc. Nat. Acad. Sci. USA* 72, 2550-2554 (1975).
- Keller, W.: Determination of the number of superhelical turns in simian virus 40 DNA by gel electrophoresis. *Proc. Nat. Acad. Sci. USA* 72, 4876-4880 (1975).
- Keller, W., U. Müller, I. Eicken, I. Wendel, and H. Zentgraf: Biochemical and ultrastructural analysis of SV40 chromatin. *Cold Spring Harbor Symp. Quant. Biol.* 42, 227-244 (1978).
- Müller, U., H. Zentgraf, I. Eicken, and W. Keller: Higher order structure of simian virus 40 chromatin. *Science* 201, 406-415 (1978).
- Dunn, A.R., M.B. Mathews, L.T. Chow, J. Sambrook, and W. Keller: A supplementary adenoviral leader sequence and its role in messenger translation. *Cell* 15, 511-526 (1978).
- Zain, S., J. Sambrook, R.J. Roberts, W. Keller, M. Fried, and A.R. Dunn: Nucleotide sequence of the leader segments in a cloned copy of adenovirus-2 fiber mRNA. *Cell* 16, 851-861 (1979).

- Küpper, H., W. Keller, C. Kurz, S. Forss, H. Schaller, R. Franze, K. Strohmaier, O. Marquardt, V.G. Zaslavsky, and P.H. Hofschneider: Cloning of cDNA of major antigen of foot and mouth disease virus and expression in E.coli. *Nature* 289, 555-559 (1981).
- Weingärtner, B. and W. Keller: Transcription and processing of adenoviral RNA by extracts from HeLa cells. *Proc. Nat. Acad. Sci. USA* 78, 4092-4096 (1981).
- Hernandez, N. and W. Keller: Splicing of *in vitro* synthesized messenger RNA precursors in HeLa cell extracts. *Cell* 35, 89-99 (1983).
- Krämer, A., W. Keller, B. Appel, and R. Lührmann: The 5' terminus of the RNA moiety of U1 small nuclear ribonucleoprotein particles is required for the splicing of messenger RNA precursors. *Cell* 38, 299-307 (1984).
- Sergeant, A., D. Bohmann, H. Zentgraf, H. Weiher, and W. Keller: A transcription enhancer acts *in vitro* over distances of hundreds of basepairs on both circular and linear templates but not on chromatin-reconstituted DNA. *J. Mol. Biol.* 180, 577-600 (1984).
- Frendewey, D. and W. Keller: Stepwise assembly of a pre-mRNA splicing complex requires U-snRNPs and specific intron sequences. *Cell* 42, 355-367 (1985).
- Krämer, A. and W. Keller: Purification of a protein required for the splicing of pre-mRNA and its separation from the lariat debranching enzyme. *EMBO J.* 4, 3571-3581 (1985).
- Gick, O., A.Krämer, W. Keller, and M.L. Birnstiel: Generation of histone mRNA 3' ends by endonucleolytic cleavage of the pre-mRNA in a snRNP-dependent *in vitro* reaction. *EMBO J.* 5, 1319-1326 (1986).
- Bohmann, D., W. Keller, T. Dale, H.R. Schöler, G. Tebb, and I.W. Mattaj: A transcription factor which binds to the enhancers of SV40, immunoglobulin heavy chain, and U2 snRNA genes. *Nature* 325, 268-272 (1987).
- Christofori, G., D. Frendewey, and W. Keller: Two spliceosomes can form simultaneously and independently on synthetic double-intron messenger RNA precursors. *EMBO J.* 6, 1747-1755 (1987).
- Krämer, A., M. Frick, and W. Keller: Separation of multiple components of HeLa cell nuclear extracts required for pre-messenger RNA splicing. *J. Biol. Chem.* 262, 17630-17640 (1987).

- Humphrey, T., G. Christofori, V. Lucijanac, and W. Keller: Cleavage and polyadenylation of messenger RNA precursors *in vitro* occurs within large and specific 3' processing complexes. *EMBO J.* 6, 4159-4168 (1987).
- Friendewey, D., A. Krämer, and W. Keller: Different small nuclear ribonucleoprotein particles are involved in different steps of splicing complex formation. *Cold Spring Harbor Symp. Quant. Biol.* 52, 287-298 (1987).
- Christofori, G. and W. Keller: 3' cleavage and polyadenylation of mRNA precursors *in vitro* requires a poly(A) polymerase, a cleavage factor, and a snRNP. *Cell* 54, 875-889 (1988).
- Christofori, G. and W. Keller: Poly(A) polymerase purified from HeLa cell nuclear extract is required for both cleavage and polyadenylation of pre mRNA *in vitro*. *Mol. Cell. Biol.* 9, 193-203 (1989).
- Lang, K.M. and W. Keller: Sequence requirements in different steps of the pre-mRNA splicing reaction: Analysis by the RNA modification-exclusion technique. *Mol. Cell. Biol.* 10, 4942-4947 (1990).
- Bardwell, V.J., M. Wickens, S. Bienroth, W. Keller, B.S. Sproat, and A.I. Lamond: Site-directed ribose methylation identifies 2'-OH groups in polyadenylation substrates critical for AAUAAA recognition and poly(A) addition. *Cell* 65, 125-133 (1991).
- Lingner, J., I. Radtke, E. Wahle, and W. Keller: Purification and characterization of poly(A) polymerase from *Saccharomyces cerevisiae*. *J. Biol. Chem.* 266, 8741-8746 (1991).
- Suter-Crazzolara, C. and W. Keller: Organization and transient expression of the gene for human U11 snRNA. *Gene Expression* 1, 91-102 (1991).
- Bienroth, S., E. Wahle, C. Suter-Crazzolara, and W. Keller: Purification of the cleavage and polyadenylation factor involved in the 3'-processing of messenger RNA precursors. *J. Biol. Chem.* 266, 19768-19776 (1991).
- Keller, W., S. Bienroth, K.M. Lang, and G. Christofori: Cleavage and polyadenylation factor CPF specifically interacts with the pre-mRNA 3' processing signal AAUAAA. *EMBO J.* 10, 4241-4249 (1991).
- Wahle, E., Martin, G., Schiltz, E., and W. Keller: Isolation and expression of cDNA clones encoding mammalian poly(A) polymerase. *EMBO J.* 10, 4251-4257 (1991).

- Lingner, J., Kellermann, J., and W. Keller: Cloning and expression of the essential gene for poly(A) polymerase from *S. cerevisiae*. *Nature* 354, 496-498 (1991).
- Bienroth, S., Keller, W., and E. Wahle: Assembly of a processive messenger RNA polyadenylation complex. *EMBO J.* 12, 585-594 (1993).
- Lingner, J. and W. Keller: 3'-end labelling of RNA with recombinant yeast poly(A) polymerase. *Nucleic Acids Res.* 21, 2917-2920 (1993).
- Gunderson, S., K. Beyer, G. Martin, W. Keller, W.C. Boelens, and I.W. Mattaj: The human U1A snRNP protein regulates polyadenylation via a direct interaction with poly(A) polymerase. *Cell* 76, 531-541 (1994).
- O'Connell, M.A. and W. Keller: Purification and properties of double-stranded RNA-specific adenosine deaminase. *Proc. Nat. Acad. Sci. USA* 91, 10596-10600 (1994).
- Jenny, A. H.-P. Hauri, and W. Keller: Characterization of cleavage and polyadenylation specificity factor and cloning of its 100-Kilodalton subunit. *Mol. Cell. Biol.* 14, 8183-8190 (1994).
- Minvielle-Sebastia, L., P.J. Preker, and W. Keller: RNA14 and RNA15 proteins as components of a yeast pre-mRNA 3'-end processing factor. *Science* 266, 1702-1705 (1994).
- O'Connell, M.A., S. Krause, M. Higuchi, J.J. Hsuan, N.F. Totty, A. Jenny, and W. Keller: Cloning of cDNAs encoding mammalian double-stranded RNA-specific adenosine deaminase. *Mol. Cell. Biol.* 15, 1389-1397 (1995).
- Melcher, T., S. Maas, M. Higuchi, W. Keller, and P.H. Seeburg: Editing of AMPA receptor GluR-B pre-mRNA *in vitro* reveals site-selective adenosine to inosine conversion. *J. Biol. Chem.* 270, 8566-8570 (1995).
- Preker, P.J., J.Lingner, L. Minvielle-Sebastia, and W. Keller: The *FIP1* gene encodes a component of a yeast pre-mRNA polyadenylation factor that directly interacts with poly(A) polymerase. *Cell* 81, 379-389 (1995).
- Jenny, A., and W. Keller: Cloning of cDNAs encoding the 160 kDa subunit of the bovine cleavage and polyadenylation specificity factor. *Nucleic Acids Res.* 14, 2629-2635 (1995).

- Maas, S., T. Melcher, A. Herb, P.H. Seeburg, W. Keller, S. Krause, M. Higuchi, and M.A. O'Connell: Structural requirements for RNA editing in glutamate receptor pre-mRNAs by recombinant double-stranded RNA adenosine deaminase. *J. Biol. Chem.* 271, 12221-12226 (1996).
- Rüegsegger, U., K. Beyer, and W. Keller: Purification and characterization of human cleavage factor I<sub>m</sub> involved in the 3' end processing of messenger RNA precursors. *J. Biol. Chem.* 271, 6107-6113 (1996).
- Martin, G. and W. Keller: Mutational analysis of mammalian poly(A) polymerase identifies a region for primer binding and a catalytic domain, homologous to the family X polymerases, and to other nucleotidyltransferases. *EMBO J.* 15, 2593-2603 (1996).
- Ohnacker, M., L. Minvielle-Sebastia, and W. Keller: The *Schizosaccharomyces pombe pla1* gene encodes a poly(A) polymerase and can functionally replace its *Saccharomyces cerevisiae* homologue. *Nucleic Acids Res.* 24, 2585-2591 (1996).
- Jenny, A., L. Minvielle-Sebastia, P.J. Preker, and W. Keller: Sequence similarity between the 73-kDa protein of mammalian CPSF and a subunit of yeast polyadenylation factor I. *Science* 254, 1514-1517 (1996).
- O'Connell, M.A., A. Gerber, and W. Keller: Purification of human double-stranded RNA-specific editase 1 (hRED1) involved in editing of brain glutamate receptor B pre-mRNA. *J. Biol. Chem.* 272, 473-478 (1997).
- Gerber, A., M.A. O'Connell, and W. Keller: Two forms of human double-stranded RNA-specific editase 1 (hRED1) generated by the insertion of an Alu cassette. *RNA* 3, 453-463 (1997).
- Barabino, S.M.L., W. Hübner, A. Jenny, L. Minvielle-Sebastia, and W. Keller: The 30 kDa subunit of mammalian cleavage and polyadenylation specificity factor and its yeast homolog are RNA binding zinc finger proteins. *Genes Dev.* 11, 1703-1716 (1997).
- Minvielle-Sebastia, L., P.J. Preker, T. Wiederkehr, Y. Strahm, and W. Keller: The major yeast poly(A)-binding protein is associated with cleavage factor IA and functions in pre-messenger RNA 3'-end formation. *Proc. Nat. Acad. Sci. USA.* 94, 7897-7902 (1997).
- Preker, P.J., M. Ohnacker, L. Minvielle-Sebastia, and W. Keller: A multisubunit 3'-end processing factor from yeast containing poly(A) polymerase and homologues of the subunits of mammalian cleavage and polyadenylation specificity factor. *EMBO J.* 16, 4727-4737 (1997).

- Beyer, K., T. Dandekar, and W. Keller: RNA-ligands selected by cleavage stimulation factor contain distinct sequence motifs that function as downstream elements in 3'-end processing of pre-mRNA. *J. Biol. Chem.* 272, 26769-26779 (1997).
- Preker, P.J. and W. Keller: The HAT helix, a repetitive motif implicated in RNA processing. *Trends Biochem. Sci.* 23, 15-16 (1998).
- Rüegsegger, U., D. Blank, and W. Keller: Human pre-mRNA cleavage factor  $I_m$  is related to spliceosomal SR proteins and can be reconstituted in vitro from recombinant subunits. *Mol. Cell* 1, 243-253 (1998).
- Martin, G. and W. Keller: Tailing and 3'-end labeling of RNA with yeast poly(A) polymerase and various nucleotides. *RNA* 4, 226-230 (1998).
- Birse, C.E., L. Minvielle-Sebastia, B.A. Lee, W. Keller, and N.J. Proudfoot: Coupling termination of transcription to messenger RNA maturation in yeast. *Science* 280, 298-301 (1998).
- Dandekar, T., K. Beyer, P. Bork, M.-R. Kenealy, K. Pantopoulos, M.W. Hentze, V. Sonntag-Buck, G. Flouriou, F. Gannon, W. Keller, and S. Schreiber: Systematic genomic screening and analysis of mRNA in the untranslated region: combining computational approaches and experiment. *Bioinformatics* 14, 271-278 (1998).
- Nemeroff, M.E., S.M.L. Barabino, Y. Li, W. Keller, and R.M. Krug: Influenza virus NS1 protein interacts with the cellular 30 KD subunit of CPSF and inhibits 3' end formation of cellular pre-mRNAs. *Mol. Cell* 1, 991-1000 (1998).
- Gerber, A., H. Grosjean, T. Melcher, and W. Keller: Tad1p, a yeast tRNA-specific adenosine deaminase, is related to the mammalian pre-mRNA editing enzymes ADAR1 and ADAR2. *EMBO J.* 17, 4780-4789 (1998).
- Minvielle-Sebastia, L., K. Beyer, A.M. Krecic, R.E. Hector, M.S. Swanson, and W. Keller: Control of cleavage site selection during mRNA 3'-end formation by a yeast hnRNP. *EMBO J.* 17, 7454-7468 (1998).
- Gerber, A.P. and W. Keller: An adenosine deaminase that generates inosine at the wobble position of tRNAs. *Science* 286, 1146-1149 (1999).

- Martin, G., P. Jenö, and W. Keller: Mapping of ATP binding regions in poly(A) polymerases by photoaffinity labeling and by mutational analysis identifies a domain conserved in many nucleotidyltransferases. *Protein Science* 8, 2380-2391 (1999).
- Ohnacker, M., S.M.L. Barabino, P.J. Preker, and W. Keller: The WD-repeat protein Pfs2p bridges two essential factors within the yeast pre-mRNA 3'-end processing complex. *EMBO J.* 19, 37-47 (2000).
- Keegan, L.P., A.P. Gerber, J. Brindle, R. Leemans, A. Gallo, W. Keller, and M.A. O'Connell: The properties of a tRNA-specific adenosine deaminase from *Drosophila melanogaster* support an evolutionary link between pre-mRNA editing and tRNA modification. *Mol. Cell. Biol.* 20, 825-833 (2000).
- Vagner, S., U. Rügsegger, S.I. Gunderson, W. Keller, and I.W. Mattaj: Position-dependent inhibition of the cleavage step of pre-mRNA 3'-end processing by U1 snRNP. *RNA* 6, 178-188 (2000).
- Barabino, S.M.L., M. Ohnacker, and W. Keller: Distinct roles of two Yth1p domains in 3'-end cleavage and polyadenylation of yeast pre-mRNAs. *EMBO J.* 19, 3778-3787 (2000).
- Martin, G., W. Keller, and S. Doublé: Crystal structure of mammalian poly(A) polymerase in complex with an analog of ATP. *EMBO J.* 19, 4193-4203 (2000).
- De Vries, H., U. Rügsegger, W. Hübner, A. Friedlein, H. Langen, and W. Keller: Human pre-mRNA cleavage factor  $Il_m$  contains homologs of yeast proteins and bridges two other cleavage factors. *EMBO J.* 19, 5895-5904 (2000).
- Fahrenkrog, B., W. Hübner, A. Mandinova, N. Panté, W. Keller, and U. Aebi: The yeast nucleoporin Nup53p specifically interacts with Nic96p and is directly involved in nuclear protein import. *Mol. Biol. Cell* 11, 3885-3896 (2000).
- Dichtl, B. and W. Keller: Recognition of polyadenylation sites in yeast pre-mRNAs by cleavage and polyadenylation factor. *EMBO J.* 20, 3197-3209 (2001).
- Morlando, M., P. Greco, B. Dichtl, A. Fatica, W. Keller, and I. Bozzoni: Functional analysis of yeast snRNA 3'-end formation mediated by uncoupling of cleavage and polyadenylation. *Mol. Cell. Biol.* 22, 1379-1389 (2002).



- Dichtl, B., D. Blank, M. Sadowski, W. Hübner, S. Weiser, and W. Keller: Yhh1p/Cft1p directly links poly(A) site recognition and RNA polymerase II transcription termination. *EMBO J.* 21, 4125-4135 (2002).
- Dichtl, B., D. Blank, M. Ohnacker, A. Friedlein, D. Roeder, H. Langen, and W. Keller: A role for *SSU72* in balancing RNA polymerase II transcription elongation and termination. *Mol. Cell* 10, 1139-1150 (2002).
- Wolf, J., A.P. Gerber, and W. Keller: *tadA*, an essential tRNA-specific adenosine deaminase from *Escherichia coli*. *EMBO J.* 21, 3841-3851 (2002).
- Sadowski, M., B. Dichtl, W. Hübner, and W. Keller: Independent functions of yeast Pcf11p in pre-mRNA 3' end processing and in transcription termination. *EMBO J.* 22, 2167-2177 (2003).
- Kyburz, A., M. Sadowski, B. Dichtl, and W. Keller: The role of the yeast cleavage and polyadenylation factor subunit Ydh1p/Cft2p in pre-mRNA 3'-end formation. *Nucleic Acids Res.* 31, 3936-3945 (2003).
- Kaufmann, I., G. Martin, A. Friedlein, H. Langen, and W. Keller: Human Fip1 is a subunit of CPSF that binds to U-rich RNA elements and stimulates poly(A) polymerase. *EMBO J.* 23, 616-626 (2004).
- Martin, G. and W. Keller: Sequence motifs that distinguish ATP(CTP):tRNA nucleotidyl transferases from eubacterial poly(A) polymerases. *RNA* 10, 899-906 (2004).
- Dichtl, B., R. Aasland, and W. Keller: Functions for *S. cerevisiae* Swd2p in 3' end formation of specific mRNAs and snoRNAs and global histone 3 lysine 4 methylation. *RNA* 10, 965-977 (2004).
- Dettwiler, S., C. Aringhieri, S. Cardinale, W. Keller, and S.M.L. Barabino: Distinct sequence motifs within the 68-kDa subunit of cleavage factor I mediate RNA binding, protein-protein interactions and subcellular localization. *J. Biol. Chem.* 279, 35788-35797 (2004).
- Martin, G., A. Möglich, W. Keller, and S. Doublé: Biochemical and structural insights into substrate binding and catalytic mechanism of mammalian poly(A) polymerase. *J. Mol. Biol.* 341, 911-925 (2004).
- Vanacova, S., J. Wolf, G. Martin, D. Blank, S. Dettwiler, A. Friedlein, H. Langen, G. Keith, and W. Keller: A new yeast poly(A) polymerase involved in RNA quality control. *PLoS Biology* 3, 987-997 (2005) [e189].
- Ingham, R.J., K. Colwill, C. Howard, S. Dettwiler, C.S.H. Lim, J. Yu, K. Hersi, G. Gish, G. Mbamalu, L.

- Taylor, B. Yeung, G. Vassilovski, M. Amin, F. Chen, L. Matskova, G. Winberg, I. Ernberg, R. Linding, P. O'Donnell, A. Starostine, W. Keller, P. Metalnikov, C. Stark, and T. Pawson: WW domains provide a platform for the assembly of multi-protein networks. *Mol. Cell. Biol.* 25, 7092-7106 (2005).
- Kyburz, A., A. Friedlein, H. Langen, and W. Keller: Direct interactions between subunits of CPSF and the U2 snRNP contribute to the coupling of pre-mRNA 3' end processing and splicing. *Mol. Cell* 23, 195-205 (2006).
- Millevoi, S., C. Loulergue, S. Dettwiler, S.Z. Karaa, W. Keller, M. Antoniou, and S. Vagner: An interaction between U2AF65 and CF I<sub>m</sub> links the splicing and 3' end processing machineries. *EMBO J.* 25, 4854-4864 (2006).
- Danckwardt, S., I. Kaufmann, M. Gentzel, K.U. Foerstner, A.-S. Gantzert, N.H. Gehring, G. Neu-Yilik, P. Bork, W. Keller, M. Wilm, M.W. Hentze, and A.E. Kulozik: Splicing factors stimulate polyadenylation via USEs at non-canonical 3' end formation signals. *EMBO J.* 26, 2658-2669 (2007).
- Coseno, M.E., G. Martin, C. Berger, G. Gilmartin, W. Keller, and S. Doublé: Crystal structure of the 25 kDa subunit of human cleavage factor I<sub>m</sub>. *Nucleic Acids Res.* 36, 3474-3483 (2008).
- Garas, M., B. Dichtl, and W. Keller: The role of the putative 3' end processing endonuclease Ysh1p in mRNA and snoRNA synthesis. *RNA* 14, 2671-2684 (2008).
- San Paolo, S., S. Vanacova, L. Schenk, T. Scherrer, D. Blank, W. Keller, and A.P. Gerber: Distinct roles of non-canonical poly(A) polymerases in RNA metabolism. *PLoS Genetics* 5, e1000555 (2009).
- Martin, G., A. Ostareck-Lederer, A. Chari, N. Neuenkirchen, S. Dettwiler, D. Blank, U. Rügsegger, U. Fischer and W. Keller: Arginine methylation in subunits of mammalian pre-mRNA cleavage factor I. *RNA* 16, 1646-1659 (2010).
- Rammelt, C., B. Biter, M. Zavolan and W. Keller: PAPD5, a noncanonical poly(A) polymerase with an unusual RNA-binding motif. *RNA* 17, 1737-1746 (2011).
- Martin, G., A.R. Gruber, W. Keller and M. Zavolan: Genome-wide analysis of pre-mRNA 3' end processing reveals a decisive role of human cleavage factor I in the regulation of 3' UTR length. *Cell Reports* 1, 753-763 (2012).

**Other publications:**

Keller, W.: Zum Einfluss von Heparinen auf die Aktivität der Mono- und Diaminoxidase.

M.D. Thesis. Medical Academy of Düsseldorf (1962).

Goedde, H.W. and W. Keller: Metabolic pathways in maple syrup urine disease. In: "Amino acid metabolism and genetic variation", W.L. Nyhan ed. McGraw Hill, N.Y. pp. 191-214 (1967).

Goedde, H.W., W. Keller, K.G. Blume, W. Hense, and D. Brackertz: Zur Genetik und Biochemie der Ahornsirupkrankheit. (branched chain ketoaciduria). Dtsch. Med. Wschr. 93, 903-908 (1968).

Ozanne, B., A. Vogel, P.A. Sharp, W. Keller, and J. Sambrook: Transcription of SV40 DNA sequences in different transformed cell lines. In: "Possible episomes in eukaryotes", L.G. Silvestri ed. North-Holland Publishing Company (Amsterdam-London) pp. 176-182 (1973).

Keller, W. and R. Crouch: The relationship of viral to cellular ribonuclease H. In: "Possible episomes in eukaryotes", L.G. Silvestri ed. North-Holland Publishing Company (Amsterdam-London) pp. 123-129 (1973).

Keller, W.: The purification of a DNA relaxing enzyme from human tissue culture cells and its interaction with supercoiled DNA. Ph.D. Thesis, State University of New York, Stony Brook (1974).

Sambrook, J., A. Jackson, W. Keller, B. Ozanne, P.A. Sharp, B. Sugden: Transcription of SV40 in lytically infected and transformed cells. In: Kolber AR, ed. Tumor virus-host cell interaction. New York, Plenum Press, pp. 91-116 (1975).

Zentgraf, H., W. Keller, and U. Müller: The structure of SV40 chromatin. Phil. Trans. R. Soc. Lond. B. 283, 299 -203 (1978).

Keller, W.: The RNA lariat: a new ring to the splicing of mRNA precursors. Cell 39, 423-425 (1984).

Tebb, G., I.W. Mattaj, W. Keller, and D. Bohmann: Characterization of an enhancer-binding protein. In: "RNA polymerase and the regulation of transcription", W.S. Reznikoff et al., eds. Elsevier Publishing Company (Amsterdam) pp. 421-426 (1987).

Matthias, P., T. Gerster, M. Thali, D. Bohmann, W. Keller, W. Zürcher, J. Jiricny, and W. Schaffner: Determinants of cell type specificity within the immunoglobulin heavy chain gene enhancer. In: Alfred Benzon Symposium, N.D. Kjeldgaard and J. Forchhammer, eds. Vol. 24, pp. 36-47 (1987).

- Keller, W.: Posttranscriptional processing (splicing and 3'-end formation) of nuclear messenger RNA precursors *in vitro*. In: "The molecular basis of viral replication", R. Perez Bercoff, ed. NATO ASI Series A, vol. 136, pp. 99-111, Plenum Press (1987).
- Steitz, J.A., D.L. Black, V. Gerke, K.A. Parker, A. Krämer, D. Frendewey, and W. Keller: Functions of the abundant U-snRNPs. In: "Structure and function of major and minor small nuclear ribonucleoprotein particles." M.L. Birnstiel ed. Springer-Verlag, pp. 115-154 (1988).
- Krämer, A. and W. Keller: Preparation and fractionation of mammalian extracts active in pre-mRNA splicing. *Methods Enzymol.* 181, 3-19 (1990).
- Wahle, E. and W. Keller: The biochemistry of 3'-end cleavage and polyadenylation of messenger RNA precursors. *Ann. Rev. Biochem.* 61, 419-440 (1992).
- Wahle, E. and W. Keller: Polyadenylation and other mRNA modifications. In: "RNA processing: A practical approach", Oxford University Press; Vol. II, pp. 1-34 (1994).
- Keller, W.: 3'-end cleavage and polyadenylation of nuclear messenger RNA precursors. In: "Pre-mRNA processing", A.I. Lamond ed. Springer-Verlag/R.G. Landes Bioscience Publishers. pp.113-134 (1995).
- Keller, W.: No end yet to messenger RNA 3' processing! *Cell* 81, 829-832 (1995).
- Wahle, E. and W. Keller: The biochemistry of polyadenylation. *Trends Biochem. Sci.* 21, 247-250 (1996).
- Keller, W. and L. Minvielle-Sebastia: A comparison of mammalian and yeast pre-mRNA 3'-end processing. *Curr. Op. Cell Biol.* 9, 329-336 (1997).
- Wahle, E. and W. Keller: Capping, methylation, and 3'-end formation of messenger RNA precursors. In: "Eukaryotic mRNA processing" *Frontiers in Molecular Biology*, B.D. Hames and D.M. Glover eds. Oxford University Press; pp. 280-309 (1997).
- Bass, B.L., K. Nishikura, W. Keller, P.H. Seeburg, R.B. Emeson, M.A. O'Connell, C.E. Samuel, and A. Herbert: A standardized nomenclature for adenosine deaminases that act on RNA. *RNA* 3, 947-949 (1997).

- Minvielle-Sebastia, L. and W. Keller: mRNA polyadenylation and its coupling to other RNA processing reactions and to transcription. *Curr. Op. Cell Biol.* 11, 352-357 (1999).
- Keller, W., J. Wolf, and A. Gerber: Editing of messenger RNA precursors and of tRNAs by adenosine to inosine conversion. *FEBS Letters* 452, 71-76 (1999).
- Keller, W.: In the beginning, there was RNA. *Science* 285, 668-669 (1999).
- Barabino, S.M.L. and W. Keller: Last but not least: regulated poly(A) tail formation. *Cell* 99, 9-11 (1999).
- Gerber, A.P. and W. Keller: RNA editing by base deamination: more enzymes, more targets, new mysteries. *Trends Biochem. Sci.* 26, 376-384 (2001).
- Keller, W. and G. Martin: Gene expression: Reviving the message. *Nature* 419, 267-268 (2002).
- Schaub, M. and W. Keller: RNA editing by adenosine deaminases generates RNA and protein diversity. *Biochimie* 84, 791-803 (2002).
- Martin, G. and W. Keller: RNA-specific ribonucleotidyl transferases. *RNA* 13, 1834-1849 (2007).
- Martin, G. S. Doubl  , and W. Keller: Determinants of substrate specificity in RNA-dependent nucleotidyl transferases. *Biochem. Biophys. Acta* 1779, 206-216 (2008).

**Publications by members of the group on which I am not a coauthor:**

Cattaneo, R., H. Will, N. Hernandez, and H. Schaller: Signals regulating hepatitis B surface antigen transcription. *Nature* 305, 336-338 (1983).

Shaw, P.E., D. Bohmann, and A. Sergeant: The SV40 enhancer influences viral late transcription *in vitro* but not on replicating templates. *EMBO J.* 4, 3247-3252 (1985).

Brown, J.W.S., G. Feix, and D. Frendewey: Accurate *in vitro* splicing of two pre-mRNA plant introns in a HeLa cell nuclear extract. *EMBO J.* 5, 2749-2758 (1986).

Krämer, A.: Analysis of RNase-A-resistant regions of adenovirus 2 major late precursor-mRNA in splicing extracts reveals an ordered interaction of nuclear components with the substrate RNA. *J.Mol.Biol.* 196, 559-573 (1987).

Krämer, A.: Fractionation of HeLa cell nuclear extracts reveals minor small nuclear ribonucleoprotein particles. *Proc. Natl. Acad. Sci. USA* 84, 8408-8412 (1987).

Gick, O., A. Krämer, A. Vasserot, and M.L. Birnstiel: A heat-labile regulatory factor is required for 3' processing of histone pre-mRNAs. *Proc. Natl. Acad. Sci. USA* 84, 8937-8940 (1987).

Krämer, A.: Presplicing complex formation requires two proteins and U2 snRNP. *Genes Dev.* 2, 1155-1167 (1988).

Krämer, A.: Purification of mammalian small nuclear ribonucleoprotein particles active in RNA processing. *Meth. Enzymol.* 181, 215-232 (1990).

Krämer, A.: Site-specific degradation of the RNA of small nuclear ribonucleoprotein particles with complementary oligodeoxynucleotides and RNase H. *Meth. Enzymol.* 181, 284-292 (1990).

Utans, U. and A. Krämer: Splicing factor SF4 is dispensable for the formation of a functional splicing complex and participates in the subsequent splicing reaction. *EMBO J.* 9, 4119-4126 (1990).

Krämer, A., U. Utans, W. Keller, and A. Lamond: Components involved in nuclear pre-mRNA splicing. *Mol. Biol. Reports* 14, 199-200 (1990).

- Krämer, A.: Purification of mammalian small nuclear ribonucleoprotein particles active in RNA processing. *Methods Enzymol.* 181, 215-232 (1990).
- Krämer, A.: Site-specific degradation of the RNA of small nuclear ribonucleoprotein particles with complementary oligodeoxynucleotides and RNase H. *Methods Enzymol.* 181, 284-292 (1990).
- Utans, U. and A. Krämer: Splicing factor SF4 is dispensable for the formation of a functional splicing complex and participates in the subsequent splicing reaction. *EMBO J.* 9, 4119-4126 (1990).
- Wahle, E.: Purification and characterization of a mammalian polyadenylate polymerase involved in the 3' end processing of messenger RNA precursors. *J. Biol. Chem.* 266, 3131-3139 (1991).
- Wahle, E.: A novel poly(A)-binding protein acts as a specificity factor in the second phase of messenger RNA polyadenylation. *Cell* 66, 759-768 (1991).
- Krämer, A. and U. Utans: Three protein factors (SF1, SF3 and U2AF) function in presplicing complex formation in addition to snRNPs. *EMBO J.* 10, 1503-1509 (1991).
- Wahle, E.: The end of the message: 3'-end processing leading to polyadenylated RNA. *BioEssays* 14, 113-118 (1992).
- Nemeroff, M.E., U. Utans, A. Krämer, and R.M. Krug: Identification of cis-acting intron and exon regions in influenza virus NS1 mRNA that inhibit splicing and cause the formation of aberrantly sedimenting pre-splicing complexes. *Mol. Cell. Biol.* 12, 962-970 (1992).
- Utans, U., S.E. Behrens, R. Lührmann, R. Kole, and A. Krämer: A splicing factor that is inactivated during in vivo heat shock is functionally equivalent to the [U4/U6.U5] triple snRNP-specific proteins. *Genes Dev.* 6, 631-641 (1992).
- Fox, C.A., M.D. Sheets, E. Wahle, and M. Wickens: Polyadenylation of maternal mRNA during oocyte maturation: poly(A) addition in vitro requires a regulated RNA binding activity and a poly(A) polymerase. *EMBO J.* 11, 5021-5032 (1992).
- Sachs, A. and E. Wahle: Poly(A) tail metabolism and function in eucaryotes. *J. Biol. Chem.* 268, 22955-22958 (1993).
- Wahle, E., A. Lustig, P. Jenö, and P. Maurer: Mammalian poly(A)-binding protein II: physical properties and binding to polynucleotides. *J. Biol. Chem.* 268, 2937-2945 (1993).

- Krause, S., S. Fakan, K. Weis, and E. Wahle: Immunodetection of poly(A)-binding protein II in the cell nucleus. *Exp. Cell Res.* 214, 75-82 (1994).
- Wahle, E.: Poly(A) tail length control is caused by termination of processive synthesis. *J. Biol. Chem.* 270, 2800-2808 (1995).
- Wahle, E. 3'-end cleavage and polyadenylation of mRNA precursors. *Biochim. Biophys. Acta* 1261, 183-194 (1995).
- Nemeth, A., S. Krause, D. Blank, A. Jenny, P. Jenö, A. Lustig, and E. Wahle: Isolation of genomic and cDNA clones encoding bovine poly(A) binding protein II. *Nucleic Acids Res.* 23, 4034-4041 (1995).
- Schul, W., B. Groenhout, K. Koberna, Y. Takagaki, A. Jenny, E.M.M. Manders, I. Raska, R. van Driel, and L. de Jong: The RNA 3' cleavage factors CstF64kD and CPSF100kD are concentrated in nuclear domains closely associated with coiled bodies and newly synthesized RNA. *EMBO J.* 15, 2883-2892 (1996).
- O'Connell, M.A.: RNA editing: Rewriting receptors. *Curr. Biol.* 7, 437-439 (1997).
- Cunha, C., J. Monjardino, D. Chang, S. Krause, and M. Carmo-Fonseca: Localization of hepatitis delta virus RNA in the nucleus of human cells. *RNA* 4, 680-693 (1998).
- O'Connell, M.A., A. Gerber, and L.P. Keegan: Purification of native and overexpression of recombinant double-stranded RNA-specific adenosine deaminases. *Methods Companion, Methods Enzymol.* 15, 51-62 (1998).
- Wiederkehr, T., R.F. Prétot, and L. Minvielle-Sebastia: Synthetic lethal interactions with conditional poly(A) polymerase alleles identify *LCP5*, a gene involved in 18S rRNA maturation. *RNA* 4, 1357-1372 (1998).
- Wahle, E. and U. Rügsegger: 3'-end processing of pre-mRNA in eukaryotes. *FEMS Microbiology Reviews* 23, 277-295 (1999).
- Cmarko, D., P.J. Verschure, T.E. Martin, M.E. Dahmus, S. Krause, X.-D. Fu, R. van Driel, and S. Fakan: Ultrastructural analysis of transcription and splicing in the cell nucleus after BrUTP-microinjection. *Mol. Biol. Cell* 10, 211-223 (1999).



- Maas, S., A.P. Gerber, and A. Rich: Identification and characterization of a human tRNA-specific adenosine deaminase related to the ADAR family of pre-mRNA editing enzymes. *Proc. Nat. Acad. Sci. (USA)* 96, 8895-8900 (1999).
- Stuurman, N., M. Haner, B. Sasse, W. Hübner, B. Suter, and U. Aebi: Interactions between coiled-coil proteins: Drosophila lamin Dm0 binds to the bicaudal-D protein. *Eur. J. Cell Biol.* 78, 278-287 (1999).
- Paupard, M.-C., M.A. O'Connell, A.P. Gerber, and R.S. Zukin: Patterns of developmental expression of the RNA editing enzyme rADAR2. *Neuroscience* 95, 869-879 (2000).
- Hofmann, I., M. Schnölzer, I. Kaufmann, and W.W. Franke: Symplekin, a constitutive protein of karyo- and cytoplasmic particles involved in mRNA biogenesis in *Xenopus laevis* oocytes. *Mol. Biol. Cell* 13, 1665-1676 (2002).
- Morlando, M., M. Ballarino, P. Greco, E. Caffarelli, B. Dichtl, and I. Bozzoni: Coupling between snoRNP assembly and 3' processing controls box C/D snoRNA biosynthesis in yeast. *EMBO J.* 23, 2392-2401 (2004).
- Betat, H., C. Rammelt, G. Martin, and M. Mörl: Exchange of regions between bacterial poly(A) polymerase and CCA adding enzyme generates altered specificities. *Mol. Cell* 15, 389-398 (2004).
- Collart, C., J.E. Remacle, S. Barabino, L.A. van Grunsven, L. Nelles, A.Schellens, T. Van de Putte, S. Pype, D. Huylebroeck, and K. Verschueren: Smicl is a novel Smad interacting protein and cleavage and polyadenylation specificity factor associated protein. *Genes to Cells* 10, 897-906 (2005).
- Vanacova, S. and R. Stefl: The exosome and RNA quality control in the nucleus. *EMBO Rep.* 8, 651-657 (2007).